

FIRST AID TREATMENT: *Ingestion:* If the product is swallowed, do not make the person vomit. Take the person and the container to a hospital at once for medical treatment. *Skin Contact:* If you get it onto your skin, take off all contaminated clothing immediately. Wash skin at once with water, followed by soap and water. Inhalation: Unlikely route of exposure. Remove patient from exposure, keep warm and at rest. Obtain medical attention as a precaution. *Eye Contact:* After contact with eyes, immediately irrigate with eyewash solution or clean water, holding the eyelids apart, for at least 15 minutes. Obtain immediate medical attention.

NOTE TO PHYSICIAN: Gastric lavage may be effective when performed within 4 hours of ingestion. *Antidote:* Vitamin K1 (e.g. Phytomenadione or Konakion) *Dose:* Child 20mg/day, Adult 40mg/day in divided doses, orally or by injection. Monitor prothrombin time and haemoglobin levels. Maintain treatment and keep person under supervision until prothrombin times return to and remain normal. See booklet "The treatment of anticoagulant rodenticide poisoning" available from EAGLE distributors.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

STORAGE AND DISPOSAL. DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL. DO NOT STORE AND TRANSPORT NEXT TO FOOD AND ANIMAL FEED DO NOT REUSE THIS PACKAGE

STORAGE: Store in original container in a cool, dry place inaccessible to children and pets.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. KEEP AWAY FROM FOODSTUFFS

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only as indicated. For use on the farm and industrial premises (outside buildings, warehouses and stores). EAGLE® Rat and mouse bait can kill rats and mice after a single feed. Death of rodents will occur in 4-12 days. Effective against warfarin resistant rats and mice. Before laying bait inspect infested area thoroughly to determine rat and mice entry, nesting and feeding sites and water sources.

Norway rat and roof rat: Place EAGLE® wax blocks in a covered bait station to prevent access by children and domestic animals.

1. Initial bait for heavy infestations:

Place 3 - 4 EAGLE® rat and mouse wax blocks each 10m along walls or rat runs, in rat burrows and in sheltered areas where rats are known to live or feed. Bait points should be inspected and replenish every 7 days until feeding ceases.

2. Permanent bait: Place 1 block each 5-10m (or 50m² surface area) for maintaining of control.

MICE: Place 1 EAGLE® rat and mouse block per bait station every 2-4m (or 10m² surface area) along walls or runways, in nests and in sheltered areas where mice are known to live or feed. The single feeding action of EAGLE® is particularly effective against mice with their small appetites and sporadic feeding behavior.

Where necessary, for a severe infestation, place further EAGLE® at 7-14 day intervals until feeding has ceased. It is good practice to maintain a few permanent baiting points to prevent re-infestation.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.