



**7-7-7**

Keep Out Of Reach Of Children

**CAUTION**

**Balanced Nutritional Complex**

**Foliar Fertilizer**

**EVERGREEN™** is a water-soluble balanced fertilizer containing Nitrogen, Phosphorous and Potassium. It also contains micronutrients, kelp, vitamins and Humic Acid. **EVERGREEN™** can be used on a wide variety of crops such as vegetables, row and field crops, fruit and nut trees. **EVERGREEN™** promotes overall plant health and increases plant tolerance to adverse conditions.

**Guaranteed Analysis**

**F1442**

Nitrogen ( 1.5% Ammoniac Nitrogen, 5.5 % Urea) . . . . .	7 %
Available Phosphorous (P2O5) . . . . .	7 %
Soluble Potash (K2O) . . . . .	7 %
Organic solution with microelements, vitamins, kelp and Humic acid derived from: Diammonium Phosphate, Potassium, Chloride and urea. . . . .	79 %

**Net Contents: 2.5 Gallons**

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** May be harmful if swallowed, inhaled, or absorbed through the skin. May cause irritation of eyes, nose, throat, and skin. Avoid contact with eyes and skin. Avoid breathing spray mist.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance selection chart.

**WPS Uses:** Applicators and other handlers who handle this product for any use covered by the Worker Protection Standard, 40 CFR Part 170, must wear: long sleeve shirt and long pants, chemical resistant gloves (such as barrier laminate, butyl rubber 7/14 mils, nitrile rubber 7/14 mils, neoprene rubber 7/14 mils, polyvinyl chloride (PVC) 7/14 mils, or Viton 7/14 mils) and shoes with socks. For exposures in enclosed areas: a respirator with either an organic vapor-removing cartridge with a pre-filter approved for pesticides (MSHA/NIOSH approval number prefix TC-20C), or a canister approved for pesticides (MSHA/NIOSH approval prefix TC-14G) or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P, or HE prefilters.

**NON-WPS Uses:** Applicators and other handlers who handle this product for any use NOT covered by the Worker Protection Standard, 40 CFR Part 170, should wear: long-sleeved shirt and long pants, and shoes with socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if product gets inside clothing. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is not toxic to fish when used according to label instructions. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters. Do not apply, allow to drift, or drain or flush equipment onto non-target areas.

**DIRECTIONS FOR USE**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Container Disposal:** Do not re-use empty container. Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**LIMITATIONS OF LIABILITY AND REMEDIES**

The information and directions set forth on this label are believed to be reliable and should be followed carefully. Insufficient performance and damage to crops to which the product is applied may occur for a variety of reasons, including but not limited to, the failure to follow the directions on this label, the failure to adhere to good applications practice, unusual or abnormal weather conditions and the like, all of which are beyond the control of ExcelAg, Corp., Inc. (the "Company"). The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to those factors, described above, that are beyond the control of the Company. EXCEPT AS DESCRIBED IN THE PRECEDING SENTENCE, THE COMPANY MAKES NO WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, CONCERNING THE PRODUCT, INCLUDING NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR ANY PARTICULAR PURPOSE, AND NO SUCH

WARRANTY SHALL BE IMPLIED BY LAW. THE EXCLUSIVE REMEDY AGAINST THE COMPANY FOR ANY LOSSES OR DAMAGES RELATING TO THE HANDLING OR USE OF THIS PRODUCT (INCLUDING CLAIMS BASED ON CONTRACT, NEGLIGENCE, STRICT LIABILITY OR OTHER LEGAL THEORIES) SHALL BE LIMITED TO, AT THE COMPANY'S ELECTION, ONE OF THE FOLLOWING: (1) REFUND OF THE PURCHASE PRICE PAID BY BUYER OR USE FOR PRODUCT PURCHASED, OR (2) REPLACEMENT OF THE AMOUNT OF PRODUCT USED. IN NO EVENT SHALL THE COMPANY BE LIABLE FOR CONSEQUENTIAL, SPECIAL, INCIDENTAL OR INDIRECT DAMAGES OR LOSSES OF ANY NATURE, INCLUDING, BUT NOT LIMITED TO LOSS OF PROFIT OR INCOME. The Company and the seller offer this product and the buyer and user accept the same, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

**7-7-7**

Keep Out Of Reach Of Children

**CAUTION**

**FIRST AID**

**IF SWALLOWED:** Do not induce vomiting. Never give anything by mouth to an unconscious person.

**IF IN EYES:** Immediately flush eyes with plenty of water for at least 15 minutes. Seek medical attention as needed.

**IF ON SKIN:** Remove contaminated clothes. Wash exposed areas with soap and water. Seek medical attention as needed.

**IF INHALED:** Remove exposed person to fresh air. If not breathing, give artificial respiration. Get medical attention if irritation develops or persists.

**General Use Recommendations**

Ensure proper calibration of application equipment prior to product use. Verify soil and water pH to make necessary adjustments prior to defining a fertilizer program. Conduct a soil analysis to define nutritional needs to supplement. It is necessary to perform a compatibility test if other products are to be tank mixed with **EVERGREEN™**. User assumes full responsibility to ensure product compatibility. **EVERGREEN™** can be applied through irrigation systems. The injection equipment used must be in compliance with Federal, State and local regulations. It is recommended to use **EVERGREEN™** at the end of the irrigation cycle, ensuring that sufficient water run-off is allowed once the application is completed and prior to turn the system off.

**EVERGREEN™** may be irritating if in contact with the skin and eyes. Avoid inhalation of product spray. If spilled on skin or splashed in eyes, rinse affected area with running water for more than 10 minutes. Always wear gloves and goggles.

Store in closed environments and wash thoroughly after its use.

Always use **EVERGREEN™** in open, well ventilated areas.

**Mixing Instructions:** The buffering of tank water to pH 6-8 prior to the addition of **EVERGREEN™** is recommended. Add required amount of any dry flowable, wettable powder, flowable, and solution tank mix partners to partially filled tank (1/2 total volume), agitate by mechanical or hydraulic means, add **EVERGREEN™** agitate again and then add remaining EC formulation if used, and required amount of water. Continuous agitation is recommended to keep the material in proper emulsion.

**Tank Mixing Instructions:** **EVERGREEN™** may be compatible with most commonly used pesticides. If tank mixing with other materials, add products in the following order: wettable powders, dry flowables, liquid flowables, emulsifiable concentrates, and soluble materials such as fertilizers. No claim of compatibility with other products is implied. Consult the intended tank mix partner product label for appropriate application rates and use instructions. Follow the label directions for the most restrictive of label precautions and limitations. This product cannot be mixed with any product containing a label prohibition against such mixing. Read and observe the most restrictive precautionary statements and other information appearing on product labels used in mixtures. **EVERGREEN™** may be applied in conjunction with other chemically neutral liquid fertilizers. Keep the product in suspension.

**Note:** The "Directions For Use" of this product reflects the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results.

**Ground Application Instructions:** Apply material with properly calibrated hand held, mechanical or motorized spray equipment. For hand held, mechanical, or motorized applications, use 1 - 2 pints of Evergreen per acre as described on back label. Spray volume may vary and apply as a full coverage spray.

Distributed By:

Manufactured For:

**ExcelAg, Corp. U.S.A.**  
**9130 South Dadeland Blvd.**  
**Suite 1607**  
**Miami, FL 33156-7851**  
**Tel. (305) 670-0145**  
**www.excelag.com**



**7-7-7**

Keep Out Of Reach Of Children

**CAUTION**

**Balanced Nutritional Complex**

**Foliar Fertilizer**

Use higher rates at later stages

Crop Type	Rate Oz / 100 Gal	Application Timing
Cotton, sunflower	16 - 32 Oz (1 - 2 pt./acre)	<b>Seed treatment:</b> treat seeds during 20 seconds in a solution of 1 qt. <b>EVERGREEN™</b> in 10 Gal of water. <b>After planting:</b> First application at 2nd-3rd leaf stage. Second application at flower bud stage. Third application at full bloom stage.
Vegetables- direct seeding	16 - 32 Oz (1 - 2 pt./acre)	<b>Seed treatment:</b> treat seeds during 20 seconds in a solution of 1 qt. <b>EVERGREEN™</b> in 10 Gal of water. <b>After planting:</b> Start foliar sprays 15 days after germination and continue spraying with foliar sprays at 7-10 days intervals until 8-10 days prior to harvest.
Vegetables- transplant	16 - 32 Oz (1 - 2 pt./acre)	<b>Seed treatment:</b> treat seeds during 20 seconds in a solution of 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. <b>Prior to transplanting:</b> submerge seedlings for 30 seconds in a solution of 1 qt. <b>BEST™</b> , 2 Lbs. <b>SAETA™</b> and 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. <b>After transplanting:</b> Start foliar sprays 15 days after transplant and continue spraying with foliar sprays at 7-10 days intervals until 8-10 days prior to harvest.
Tomatoes, peppers, eggplant	16 - 32 Oz (1 - 2 pt./acre)	<b>Seed treatment:</b> treat seeds during 20 seconds in a solution of 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. <b>Prior to transplanting:</b> submerge seedlings for 30 seconds in a solution of 1 qt. <b>BEST™</b> , 2 Lbs. <b>SAETA™</b> and 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. <b>After transplanting:</b> Start foliar sprays 15 days after transplant (or at 2nd-4th leaf formation) and continue spraying with foliar sprays at 8-10 days intervals in tank mixes with 2 - 4 qt. /Ac. <b>BEST™</b> , in rotation with 1.7- 2.0 Lbs./Ac. <b>SAETA™</b> until 10 days prior to harvest.
Onions, Garlic	16 - 32 Oz (1 - 2 pt./acre)	<b>Seed treatment:</b> treat seeds during 20 seconds in a solution of 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. <b>Prior to transplanting:</b> submerge seedlings for 30 seconds in a solution of 1 qt. <b>BEST™</b> , 2 Lbs. <b>SAETA™</b> and 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. <b>After transplanting:</b> Start foliar sprays 15 days after transplant (at 2nd-4th leaf formation or bulb formation initiation) and continue spraying with foliar sprays at 8-10 days intervals in tank mixes with 2-4 qts./Ac. <b>BEST™</b> , in rotation with 1.7-2.0 Lbs./Ac. <b>SAETA™</b> until 10 days prior to harvest.
Strawberry	16 - 32 Oz (1 - 2 pt./acre)	<b>Seed treatment:</b> treat seeds during 20 seconds in a solution of 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. <b>Prior to transplanting:</b> submerge seedlings for 30 seconds in a solution of 1 qt. <b>BEST™</b> , 2 Lbs. <b>SAETA™</b> and 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. <b>After transplanting:</b> Start foliar sprays 15 days after transplant ( at 2nd-4th leaf formation) and continue spraying with foliar sprays at 7-10 days intervals until 8-10 days prior to harvest.
Potatoes	16 - 32 Oz (1 - 2 pt./acre)	<b>Prior to planting:</b> submerge tubers for 30 seconds in a solution of 1 qt. <b>BEST™</b> , 2 Lbs. <b>SAETA™</b> and 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. <b>After planting:</b> Start foliar sprays 15 days after transplant (or at 2nd-4th leaf formation) and continue spraying with foliar sprays at 8-10 days intervals in tank mixes with 2 qts./Ac. <b>BEST™</b> , in rotation with 2 Lbs./Ac. <b>SAETA™</b> until 10 days prior to harvest.
Beans, Soybeans, Peas, Peanuts	16 - 32 Oz (1 - 2 pt./acre)	Start applications at 2nd-3rd leaf stage, repeat applications 2 weeks later or at full bloom stage.
Wheat, Corn, Sorghum, Oat	16 - 32 Oz (1 - 2 pt./acre)	Start applications at tillering initiation (at 3rd- 4th leaf stage). Repeat application at maximum tillering stage.
Bananas, Plantains	16 - 32 Oz (1 - 2 pt./acre)	Start application at bud flower break preferably when flower bractea is exposed. Repeat application one week later and a third application 7-10 days later. Use sufficient water to ensure plant coverage.

**LIMITATIONS OF LIABILITY AND REMEDIES**

The information and directions set forth on this label are believed to be reliable and should be followed carefully. Insufficient performance and damage to crops to which the product is applied may occur for a variety of reasons, including but not limited to, the failure to follow the directions on this label, the failure to adhere to good applications practice, unusual or abnormal weather conditions and the like, all of which are beyond the control of ExcelAg, Corp., Inc. (the "Company"). The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to those factors, described above, that are beyond the control of the Company. EXCEPT AS DESCRIBED IN THE PRECEDING SENTENCE, THE COMPANY MAKES NO WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, CONCERNING THE PRODUCT, INCLUDING NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR ANY PARTICULAR PURPOSE, AND NO SUCH

WARRANTY SHALL BE IMPLIED BY LAW. THE EXCLUSIVE REMEDY AGAINST THE COMPANY FOR ANY LOSSES OR DAMAGES RELATING TO THE HANDLING OR USE OF THIS PRODUCT (INCLUDING CLAIMS BASED ON CONTRACT, NEGLIGENCE, STRICT LIABILITY OR OTHER LEGAL THEORIES) SHALL BE LIMITED TO, AT THE COMPANY'S ELECTION, ONE OF THE FOLLOWING: (1) REFUND OF THE PURCHASE PRICE PAID BY BUYER OR USE FOR PRODUCT PURCHASED, OR (2) REPLACEMENT OF THE AMOUNT OF PRODUCT USED. IN NO EVENT SHALL THE COMPANY BE LIABLE FOR CONSEQUENTIAL, SPECIAL, INCIDENTAL OR INDIRECT DAMAGES OR LOSSES OF ANY NATURE, INCLUDING, BUT NOT LIMITED TO LOSS OF PROFIT OR INCOME. The Company and the seller offer this product and the buyer and user accept the same, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

Use higher rates at later stages

Crop Type	Rate Oz / 100 Gal	Application Timing
Leafy vegetables (Lettuce, Spinach, Cauliflower, Broccoli)	16 - 32 Oz (1 - 2 pt./acre)	<b>Seed treatment:</b> treat seeds during 20 seconds in a solution of 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. <b>Prior to transplanting:</b> submerge seedlings for 30 seconds in a solution of 1 qt. <b>BEST™</b> , 2 Lbs. <b>SAETA™</b> and 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. <b>After transplanting:</b> Start foliar sprays 15 days after transplant (or at 4th-6th leaf formation) and continue spraying with foliar sprays at 8-10 days intervals in tank mixes with 2 qts./Ac. <b>BEST™</b> , in rotation with 2 Lbs./Ac. <b>SAETA™</b> until 10 days prior to harvest.
Cucurbits (watermelon, melons, squash, zucchini, etc.)	16 - 32 Oz (1 - 2 pt./acre)	<b>Seed treatment:</b> treat seeds during 20 seconds in a solution of 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. <b>Prior to transplanting:</b> submerge seedlings for 30 seconds in a solution of 1 qt. <b>BEST™</b> , 2 Lbs. <b>SAETA™</b> and 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. <b>After transplanting:</b> Start foliar sprays 7-9days after transplant (or at 2nd - 4th leaf formation) and continue spraying with foliar sprays at 8-10 days intervals in tank mixes with 2 qts./Ac. <b>BEST™</b> , in rotation with 2 Lbs./Ac. <b>SAETA™</b> until 10 days prior to harvest.
Table Grapes and Wine Grapes	16 - 32 Oz (2.0-5.0 cc/Lit)	Start applications at bud break. Repeat applications during flower formation at 8-10 days intervals.
Alfalfa, pastures	16 - 32 Oz (1 - 2 pt./acre)	Start applications 8-10 days after harvest-cut. Repeat applications at 10 day intervals.
Citrus, Avocados	16 - 32 Oz (2.0-5.0 cc/Lit)	Start applications at flowering initiation. Repeat applications at 10 day intervals. Ensure sufficient coverage on treated areas.
Rice	16 - 32 Oz (1 - 2 pt./acre)	<b>Seed treatment:</b> treat seeds during 20 seconds in a solution of 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. Continue spraying on nursery 10 days after germination and repeat 10 days later. <b>Prior to transplanting:</b> submerge seedlings for 30 seconds in a solution of 1 qt. <b>BEST™</b> , 2 Lbs. <b>SAETA™</b> and 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. <b>After transplanting:</b> Start foliar sprays 10-15 days after transplant (at 2nd/4th leaf formation or at tillering initiation) and continue spraying with foliar sprays at 8-10 days intervals in tank mixes with 2 qts./Ac. <b>BEST™</b> , in rotation with 2 Lbs./Ac. <b>SAETA™</b> until grain formation period, this is, 10 days prior to harvest. <b>To enhance grain quality:</b> apply <b>EVERGREEN™</b> in tank mix with 1 qt. <b>BEST™</b> . Repeat application 10 days later.
Asparagus	16 - 32 Oz (1 - 2 pt./acre)	<b>Seedling (crown) treatment:</b> submerge crowns for 30 seconds in a solution of 1 qt. <b>BEST™</b> , 2 Lbs. <b>SAETA™</b> and 1 qt. <b>EVERGREEN™</b> in 10 gallons of water. <b>After transplanting:</b> Start foliar sprays at 2nd - 4th leaf formation and continue spraying until flowering initiation with foliar sprays at 8-10 days intervals in tank mixes with 2 qts./Ac. <b>BEST™</b> , in rotation with 2 Lbs./Ac. <b>SAETA™</b> until grain formation period, this is, 10 days prior to harvest.
Sugar cane	48 - 80 Oz (3 - 5 pt./acre)	<b>First application:</b> Use 2 pints per acre at tillering initiation (3 weeks after planting for new plantings and 2 weeks after harvest for ratoons). <b>Second application:</b> Use 3 pints per acre at maximum tillering, or two weeks after the first application.
Okra	16 - 32 Oz (1 - 2 pt./acre)	Start applications at 3rd - 4th leaf stage, repeat applications every 7-8 days until 2 weeks prior to last harvest.
Ornamentals	16 - 32 Oz (1 - 2 pt./acre)	Start applications at 3rd - 4th leaf stage, repeat applications every 7-8 days until 2 weeks prior to last harvest.

Distributed By:

Manufactured For:

**ExcelAg, Corp. U.S.A.**  
**9130 South Dadeland Blvd.**  
**Suite 1607**  
**Miami, FL 33156-7851**  
**Tel. (305) 670-0145**  
**www.excelag.com**